

Small RNAs have a big impact on regeneration

Elizabeth J. Thatcher and James G. Patton*

Department of Biological Sciences; Vanderbilt University; Nashville, TN USA

A number of lower vertebrates including urodele amphibians and teleost fish are remarkably adept at repairing and regenerating damaged tissues and organs. Freshwater planarians are even more amazing, capable of regenerating entire body plans from small amputated fragments. In contrast, mammalian regenerative capacity is quite limited but of intense interest, especially related to human health and disease. For those organisms capable of robust regeneration, a common theme is the use of stem cells to replace complex tissues. Key questions remain as to the origin of these cells, whether there are pools of such cells that migrate to injured regions or whether they are generated on site. Beyond their origin, how are the genetic pathways that enable differentiation into multiple cell types and tissues regulated? microRNAs (miRNAs) are small noncoding RNAs that have recently been shown to play important roles in controlling stem cell self-renewal, proliferation and differentiation. Some of these are thought to be required to maintain “stemness”. Here, we summarize recent work on the role of miRNAs in stem cells and their roles during regeneration.

Small RNAs in Regeneration

miRNA biogenesis. Recently, a new class of regulatory molecules, small noncoding RNAs referred to as microRNAs (miRNAs), has been implicated in regeneration. miRNAs have been shown to regulate many cellular functions including cell specification, differentiation, proliferation and metabolism.¹⁻¹¹ A key to understanding their role is to determine how many are encoded in animal genomes

and what genes they regulate. miRNA biogenesis begins with primary transcripts synthesized by RNA polymerase II that form extended hairpin structures containing one or more miRNAs. Some of these primary transcripts originate from independent transcripts whereas others are encoded within the introns of mRNA transcripts. Primary transcripts are recognized in the nucleus by an RNase III enzyme, Drosha, which cleaves hairpin precursors for subsequent export to the cytoplasm in a Ran-GTP dependent process.^{12,13} In the cytoplasm, hairpin precursors are recognized by another RNase III enzyme, Dicer, that produces RNA duplexes (~22 nucleotide) that are incorporated into RNA Induced Silencing Complexes (RISCs).¹⁴⁻¹⁶ Either coincident or subsequent to RISC assembly, miRNAs are guided to their target mRNAs where they most often pair with 3'UTR sequences.^{17,18} Partial complementarity leads to translational repression whereas perfect complementarity leads to cleavage of the target mRNA (Fig. 1). This process is highly conserved throughout development.

A number of studies have examined the expression patterns of miRNAs during development, differentiation and regeneration. A common theme is that more and more miRNAs are expressed as differentiation proceeds.^{19,20} A smaller number are expressed in stem cells and at early stages of development, including some that are thought to be required to maintain “stemness”.²¹⁻²⁵ For animals that are adept at repairing or regenerating damaged tissues, the role of stem cells during regeneration can be simplistically boiled down to two nonexclusive mechanisms. First, a reserve of stem cells are somehow recruited to the site of damage and differentiate to restore

Key words: regeneration, microRNAs, stem cells, zebrafish

Abbreviations: miRNAs, microRNAs; RISC, RNA induced silencing complex; NGN3, neurogenin3; ES cells, embryonic stem cells

Submitted: 03/18/10

Accepted: 04/14/10

Previously published online:
www.landesbioscience.com/journals/rnabiology/article/12085

*Correspondence to: James G. Patton;
 Email: James.G.Patton@vanderbilt.edu




Figure 2. Stages of zebrafish fin regeneration. Five stages of zebrafish fin regeneration are illustrated. Initial wound healing responses apparently stimulate a number of signaling pathways that ultimately result in the formation of a blastema, a structure that contains stem cell-like cells which divide and differentiate to re-form complex tissues. miRNA expression analyses have shown that *miR-203* and *miR-133* regulate targets as part of the Fgf and Wnt cascades,^{23,48} other miRNAs are predicted to regulate downstream targets (dashed lines²³).

is elevated during early salamander tail regeneration.⁴¹ In contrast, *miR-203* must be turned off to initiate fin regeneration and allow Lef1 expression.²³ For *miR-203*, the Fuchs and Candi groups showed that it represses stem cell activity in mice by downregulating *p63*, thereby repressing ‘stemness’.^{21,25} Further identification and understanding of the role that miRNAs play in ES self-renewal and proliferation is likely to shed light on similar involvement in regeneration and vice versa (Fig. 3).

miRNAs in Planaria

Planarians are small flatworms often found in freshwater and marine environments that are capable of remarkable feats of regeneration. Numerous stem-cell like cells, called neoblasts, make up nearly 30% of the total cells in planarians and their regenerative capacity allows re-creation of whole animals, even when cut into numerous pieces.⁴²⁻⁴⁴ Using microarrays and in situ localization, a unique set of ~50–60 miRNAs were found to be highly expressed in neoblasts.^{45,46} Key protein components of miRNA effector complexes (RISC) are members of the Argonaute family. Interestingly, two genes encoding members of the Argonaute family are expressed in neoblasts and knockdown of mRNAs encoding one of these proteins (*smedwi-2*) blocked regeneration.⁴³ It is not yet clear what small RNAs are affected

by loss of *smedwi-2*, whether miRNAs or a closely related set of small RNAs referred to as piwi-associated RNAs (piRNAs),⁴² but the findings are consistent with a key role for small RNAs in controlling stem cell gene expression pathways. In this case, the loss of *smedwi-2* did not alter neoblast maintenance but affected one or more steps of differentiation. By identifying the target genes controlled by these and other small RNAs, the hope is that key genetic pathways can be identified and dissected to potentially facilitate regeneration in higher organisms and/or to possibly allow reprogramming of adult stem cells.

miRNAs Regulate Fin Regeneration

Zebrafish are able to regenerate many damaged tissues and organs including fins, hearts, retinas and spinal cords.⁴⁷ Zebrafish caudal fins are relatively simple, composed of bony rays consisting of two hemirays that create a protective shell around nerves, blood vessels and mesenchymal cells. Fin regeneration involves five steps: wound healing, mesenchymal disorganization or reorganization, blastema formation, outgrowth and termination (Fig. 2). The key is the formation of a blastema, a structure containing stem cell-like cells that are either recruited to the damaged area or originate from the de-differentiation of cells in the area. How fin

amputation sets in motion the five steps above and results in the formation of blastema cells remains largely unknown.

Recently, miRNAs have been shown to play a role in regulating fin regeneration in fish and tail regeneration in salamanders.^{23,41,48} We showed that an intact miRNA biogenesis pathway is required for caudal fin regeneration in zebrafish; inhibition of Dicer by injection of antisense morpholinos blocked regeneration.²³ Given that miRNA expression patterns change, both up and down, during regeneration, it appears that overall gene regulation requires proper control of both positive and negative regulators of regeneration. Consistent with this, some of the miRNAs that get down-regulated during regeneration, including *miR-203* and *miR-133*, are predicted to target genes that are upregulated during regeneration (*lef1*, *msxb*, *fgfr1*, *fgf20a* and *hsp60*).^{23,48} Conversely, a subset of miRNAs whose expression levels increase are predicted to target genes that are down-regulated during regeneration.²³ One example is *miR-200b* which increases during regeneration. One of its predicted targets, *bmp3*, is correspondingly down-regulated during regeneration.⁴⁹ More broadly, genes that are down-regulated during regeneration are likely involved in preserving the terminally differentiated state and are therefore normally tightly controlled, reflecting the balance




Figure 3. Stem cell maturation. During normal development and during regeneration, stem cells play key roles in self-renewal, differentiation and specification. Shown are several miRNAs that have been implicated in the maintenance and differentiation of stem cells.

between growth and possible out of control proliferation.

Another interesting aspect of recent discoveries involving miRNAs during fin regeneration is that the first two identified miRNAs regulate key players involved in Wnt and Fgf signaling.^{23,48} One of the earliest events upon damage is wound healing which results in early activation of Fgf signaling cascades, perhaps activated by hydrogen peroxide synthesis at the wound site.⁵⁰ Downstream, or coincident with Fgf signaling, a number of signaling cascades are activated including wnts, hedgehogs and TGFβ.⁵¹ One possible role for miRNA regulation is to ensure that the gradients or readouts formed by these signaling cascades are properly initiated, maintained and finally, terminated. In many ways, this may be similar to such regulation during normal development, such as the role of *miR-214* to control muscle differentiation.¹

miRNAs Regulate Muscle Regeneration

miRNAs play key roles in muscle, controlling specification, proliferation and differentiation.⁵² *miR-1* and *miR-133* are expressed in cardiac and skeletal muscle where they are transcriptionally regulated by the well known myogenic factors MyoD, myogenin, Mef2 and SRF (serum response factor).⁵³⁻⁵⁷ Expression of *miR-1* in mammals terminates the cell cycle in

cardiac and skeletal muscle progenitors and promotes terminal differentiation.^{54,57,58} *miR-133* has the opposite effect to maintain proliferating progenitor cells.⁵³ *miR-206* is expressed only in skeletal muscles and is induced by MyoD and myogenin to promote muscle differentiation.^{59,60} This is consistent with the notion that specific subsets of miRNAs are expressed that can act in opposition to either promote or repress renewal versus differentiation (Fig. 3).

Pax3 and Pax7 are known to be expressed in skeletal muscle stem cells and represent a self-renewing population of cells that are critical for muscle growth and to maintain satellite cells that are required for the regeneration of adult skeletal muscle.⁶¹ Pax3 is essential for maintenance of these cells and its migration to sites of myogenic differentiation and is regulated by *miR-27b* to rapidly initiate terminal muscle differentiation.⁶²

miRNAs in Pancreas/β-cell Regeneration

Three major pancreatic transcription factors are necessary for early development, FoxA2, Hes1 and Pdx1.⁶³⁻⁶⁵ Conditional Dicer deletion in pancreatic progenitor cells adversely affects the development of β-, δ-, exocrine and duct cells, indicating that proper miRNA-mediated gene regulation is critical for normal pancreas development, glucose metabolism and

insulin secretion.⁶⁶ Consistent with this, *miR-124a* downregulates *foxa2* expression which in turn regulates the targets of FoxA2 including *pdx1*, *kr6.2*, *sur-1*.⁶⁴

Several groups have shown that NGN3 is a “master-controller” of pancreas development.^{63,67-70} Hairy/enhancer of split (*hes1*), a notch signaling effector, tightly controls the number of Neurogenin3 (NGN-3) producing cells⁶⁵ and knockout of *hes1* can lead to ectopic pancreas development.^{63,67-70} miRNA profiling of the regenerating mouse pancreas has indicated that *miR-15a*, *-15b*, *-16* and *-195* negatively regulate translation of *ngn3* (Fig. 3).⁷¹ *miR-23* downregulates *hes1* during neuronal development and it may have a similar role in the pancreas.⁷² In the regenerating pancreas, NGN3 levels are low, despite the fact that a number of activating transcription factors upstream of NGN3 are expressed. One possibility is that miRNA control of *ngn3* limits its expression. Interestingly, NGN3 is required to regenerate hormone-producing cells and it is possible that understanding how to alter NGN3 levels might help to treat diabetes through promoting regeneration of insulin-secreting cells.

Recent studies have shown a requirement for notch signaling in neuronal stem cell maintenance in embryonic and adult mice.⁷³ *Hes1;Hes3;Hes5* triple-mutant mice lack neural stem cell populations in most of the central nervous system except

the telencephalon.⁷⁴ All other neuronal progenitor pools prematurely differentiated into neurons. In order to more completely deplete Notch signaling in the adult mouse brain, the Kageyama group knock-downed Rbpj in mice. The Notch intracellular domain (NICD) forms a complex with Rbpj to induce the expression of transcriptional repressors like Hes1.⁷³ Without repression of neuronal genes, cells prematurely differentiate. Thus, proper Notch signaling is crucial to regulate and maintain stem cell populations. miRNAs are likely to play a role in properly regulating Notch signaling, consistent with the finding that a number of miRNAs exhibit altered expression patterns upon inhibition of Notch.¹⁹

Initiation and Termination of Regeneration

The parallels to regeneration and cancer or tumor formation are many. As an example, during fin regeneration, genes that maintain the differentiated state must be turned off while genes that promote proliferation and outgrowth must be turned on. At the end of regeneration, the regulatory switches need to be reversed. It is remarkable that zebrafish fin regeneration terminates at precisely the right size, despite the fact that fish grow throughout their lifetimes. This raises the question as to what mechanisms are in place to properly terminate regeneration. We have shown that continued repression of *miR-203* during fin regeneration leads to fin overgrowth.²³ It remains unknown what signals regulate the transcriptional activity of *miR-203* but it is clear that specific miRNAs are needed for both initiation and termination. For unregulated growth in cancer, the similarities are striking. Overexpression of *miR-34* induces apoptosis and inhibits cell growth and tumor invasion in the pancreas via targeting of *bcl-2* (B-cell CLL/lymphoma-2) and *Notch1/2*.⁷⁵⁻⁷⁷ In pancreatic cancer, the action of *miR-34* must be downregulated to escape such growth controls.

Conclusion

The processes and mechanisms that allow certain organisms to undergo regeneration

of complex tissues are only just beginning to be understood. The key lies in understanding those mechanisms (Fig. 3) and especially the role that stem cells play in these processes. As more and more is learned about regeneration, key regulatory steps that control stem cells and regeneration will be uncovered opening the door to application in humans to repair injured or damaged tissues.

References

- Flynt A, Li N, Thatcher E, Solnica-Krezel L, Patton J. Zebrafish miR-214 modulates Hedgehog signaling to specify muscle cell fate. *Nat Genet* 2007; 39:259-63.
- Flynt A, Thatcher E, Burkewitz K, Li N, Liu Y, Patton J. miR-8 microRNAs regulate the response to osmotic stress in zebrafish embryos. *J Cell Biol* 2009; 185:115-27.
- Esau C, Davis S, Murray SF, Yu XX, Pandey SK, Pear M, et al. miR-122 regulation of lipid metabolism revealed by in vivo antisense targeting. *Cell Metab* 2006; 3:87-98.
- Plasterk RH. Micro RNAs in animal development. *Cell* 2006; 124:877-81.
- Bartel D. MicroRNAs: genomics, biogenesis, mechanism and function. *Cell* 2004; 116:281-97.
- Hobert O. Common logic of transcription factor and microRNA action. *Trends Biochem Sci* 2004; 29:462-8.
- Carthew RW. Gene regulation by microRNAs. *Curr Opin Genet Dev* 2006; 16:203-8.
- Hatfield SD, Shcherbata HR, Fischer KA, Nakahara K, Carthew RW, Ruohola-Baker H. Stem cell division is regulated by the microRNA pathway. *Nature* 2005; 435:974.
- Slack FJ, Weidhaas JB. MicroRNAs as a potential magic bullet in cancer. *Future Oncol* 2006; 2:73-82.
- O'Donnell KA, Wentzel EA, Zeller KI, Dang CV, Mendell JT. c-Myc-regulated microRNAs modulate E2F1 expression. *Nature* 2005; 435:839-43.
- Voorhoeve PM, le Sage C, Schrier M, Gillis AJ, Stoop H, Nagel R, et al. A genetic screen implicates miRNA-372 and miRNA-373 as oncogenes in testicular germ cell tumors. *Cell* 2006; 124:1169-81.
- Lund E, Guttinger S, Calado A, Dahlberg JE, Kutay U. Nuclear export of microRNA precursors. *Science* 2004; 303:95-8.
- Yi R, Qin Y, Macara IG, Cullen BR. Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes Dev* 2003; 17:3011-6.
- Bernstein E, Ahmet M, Denli, Gregory J, Hannon. The rest is silence. *RNA* 2001; 7:1509-21.
- Bernstein E, Caudy AA, Hammond SM, Hannon GJ. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* 2001; 409:363-6.
- Zamore P, Tuschl T, Sharp P, Bartel D. RNAi: double-stranded RNA directs the ATP-dependent cleavage of mRNA at 21 to 23 nucleotide intervals. *Cell* 2000; 101:25-33.
- Gu S, Rossi JJ. Uncoupling of RNAi from active translation in mammalian cells. *RNA* 2005; 11:38-44.
- Sen GL, Blau HM. Argonaute 2/RISC resides in sites of mammalian mRNA decay known as cytoplasmic bodies. *Nat Cell Biol* 2005; 7:633.
- Thatcher E, Flynt A, Li N, Patton J. MiRNA expression analysis during normal zebrafish development and following inhibition of the Hedgehog and Notch signaling pathways. *Dev Dyn* 2007; 236:2172-80.
- Wienholds E, Kloosterman WP, Miska E, Alvarez-Saavedra E, Berezikov E, de Bruijn E, et al. MicroRNA Expression in Zebrafish Embryonic Development. *Science* 2005; 309:310-1.
- Lena AM, Shalom-Feuerstein R, Rivetti di Val Cervo P, Aberdam D, Knight RA, Melino G, et al. miR-203 represses 'stemness' by repressing DeltaNp63. *Cell Death Differ* 2008; 15:1187-95.
- Melton C, Judson RL, Belloch R. Opposing microRNA families regulate self-renewal in mouse embryonic stem cells. *Nature* 2010; 1-8.
- Thatcher E, Paydar I, Anderson KK, Patton J. Regulation of zebrafish fin regeneration by microRNAs. *Proc Natl Acad Sci USA* 2008; 105:18384-9.
- Wang Y, Medvid R, Melton C, Jaenisch R, Belloch R. DGC8 is essential for microRNA biogenesis and silencing of embryonic stem cell self-renewal. *Nat Genet* 2007; 39:380-5.
- Yi R, Poy MN, Stoffel M, Fuchs E. A skin microRNA promotes differentiation by repressing 'stemness'. *Nature* 2008; 452:225-9.
- Houbaviy HB, Murray MF, Sharp PA. Embryonic stem cell-specific MicroRNAs. *Dev Cell* 2003; 5:351-8.
- Tang F, Hajkova P, Barton SC, Lao K, Surani MA. MicroRNA expression profiling of single whole embryonic stem cells. *Nucleic Acids Res* 2006; 34:9.
- Forstemann K, Tomari Y, Du T, Vagin VV, Denli AM, Bratu DP, et al. Normal microRNA maturation and germ-line stem cell maintenance requires Loquacious, a double-stranded RNA-binding domain protein. *PLoS Biol* 2005; 3:236.
- Bernstein E, Kim SY, Carmell MA, Murchison EP, Alcorn H, Li MZ, et al. Dicer is essential for mouse development. *Nat Genet* 2003; 35:215-7.
- Yang WJ, Yang DD, Na S, Sandusky GE, Zhang Q, Zhao G. Dicer is required for embryonic angiogenesis during mouse development. *J Biol Chem* 2005; 280:9330-5.
- Murchison EP, Partridge JF, Tam OH, Cheloufi S, Hannon GJ. Characterization of Dicer-deficient murine embryonic stem cells. *Proc Natl Acad Sci USA* 2005; 102:12135-40.
- Wang Y, Baskerville S, Shenoy A, Babiarz JE, Baehner L, Belloch R. Embryonic stem cell-specific microRNAs regulate the G₂-S transition and promote rapid proliferation. *Nat Genet* 2008; 40:1478-83.
- Giraldez AJ, Cinalli RM, Glasner ME, Enright AJ, Thomson JM, Baskerville S, et al. MicroRNAs Regulate Brain Morphogenesis in Zebrafish. *Science* 2005; 308:833-8.
- Wienholds E, Plasterk RH. MicroRNA function in animal development. *FEBS Lett* 2005; 579:5911-22.
- Heo I, Joo C, Cho J, Ha M, Han J, Kim VN. Lin28 mediates the terminal uridylation of let-7 precursor MicroRNA. *Mol Cell* 2008; 32:276-84.
- Lehrbach NJ, Armisen J, Lightfoot HL, Murfitt KJ, Bugaut A, Balasubramanian S, et al. LIN-28 and the poly(U) polymerase PUP-2 regulate let-7 microRNA processing in *Caenorhabditis elegans*. *Nat Struct Mol Biol* 2009; 16:1016-20.
- Newman MA, Thomson JM, Hammond SM. Lin-28 interaction with the Let-7 precursor loop mediates regulated microRNA processing. *RNA* 2008; 14:1539-49.
- Rybak A, Fuchs H, Smirnova L, Brandt C, Pohl EE, Nitsch R, et al. A feedback loop comprising lin-28 and let-7 controls pre-let-7 maturation during neural stem-cell commitment. *Nat Cell Biol* 2008; 10:987-93.
- Viswanathan SR, Daley GQ, Gregory RI. Selective blockade of microRNA processing by Lin28. *Science* 2008; 320:97-100.
- Thomson JM, Parker JS, Hammond SM. Microarray Analysis of miRNA Gene Expression. *Methods Enzymol* 2007; 427:107-22.

41. Sehm T, Sachse C, Frenzel C, Echeverri K. miR-196 is an essential early-stage regulator of tail regeneration, upstream of key spinal cord patterning events. *Dev Biol* 2009; 334:468-80.
42. Palakodeti D, Smielewska M, Lu YC, Yeo GW, Graveley BR. The PIWI proteins SMEDWI-2 and SMEDWI-3 are required for stem cell function and piRNA expression in planarians. *RNA* 2008; 14:1174-86.
43. Reddien PW, Oviedo NJ, Jennings JR, Jenkin JC, Sanchez Alvarado A. SMEDWI-2 is a PIWI-like protein that regulates planarian stem cells. *Science* 2005; 310:1327-30.
44. Guo T, Peters AH, Newmark PA. A Bruno-like gene is required for stem cell maintenance in planarians. *Dev Cell* 2006; 11:159-69.
45. Friedlander MR, Adamidi C, Han T, Lebedeva S, Isenbarger TA, Hirst M, et al. High-resolution profiling and discovery of planarian small RNAs. *Proc Natl Acad Sci USA* 2009; 106:11546-51.
46. Gonzalez-Estevéz C, Arseni V, Thambyrajah RS, Felix DA, Aboobaker AA. Diverse miRNA spatial expression patterns suggest important roles in homeostasis and regeneration in planarians. *Int J Dev Biol* 2009; 53:493-505.
47. Stoick-Cooper CL, Moon RT, Weidinger G. Advances in signaling in vertebrate regeneration as a prelude to regenerative medicine. *Genes Dev* 2007; 21:1292-315.
48. Yin VP, Thomson JM, Thummel R, Hyde DR, Hammond SM, Poss KD. Fgf-dependent depletion of microRNA-133 promotes appendage regeneration in zebrafish. *Genes Dev* 2008; 22:728-33.
49. Schebesta M, Lien CL, Engel FB, Keating MT. Transcriptional profiling of caudal fin regeneration in zebrafish. *Scientific World Journal* 2006; 6:38-54.
50. Niethammer P, Grabher C, Look AT, Mitchison TJ. A tissue-scale gradient of hydrogen peroxide mediates rapid wound detection in zebrafish. *Nature* 2009; 459:996-9.
51. Iovine MK. Conserved mechanisms regulate outgrowth in zebrafish fins. *Nat Chem Biol* 2007; 3:613-8.
52. Williams AH, Liu N, van Rooij E, Olson EN. MicroRNA control of muscle development and disease. *Curr Opin Cell Biol* 2009; 21:461-9.
53. Chen JF, Mandel EM, Thomson JM, Wu Q, Callis TE, Hammond SM, et al. The role of microRNA-1 and microRNA-133 in skeletal muscle proliferation and differentiation. *Nat Genet* 2006; 38:228-33.
54. Kwon C, Han Z, Olson EN, Srivastava D. MicroRNA1 influences cardiac differentiation in *Drosophila* and regulates Notch signaling. *Proc Natl Acad Sci USA* 2005; 102:18986-91.
55. Rao PK, Kumar RM, Farkhondeh M, Baskerville S, Lodish HF. Myogenic factors that regulate expression of muscle-specific microRNAs. *Proc Natl Acad Sci USA* 2006; 103:8721-6.
56. Sokol NS, Ambros V. Mesodermally expressed *Drosophila* microRNA-1 is regulated by Twist and is required in muscles during larval growth. *Genes Dev* 2005; 19:2343-54.
57. Zhao Y, Samal E, Srivastava D. Serum response factor regulates a muscle-specific microRNA that targets Hand2 during cardiogenesis. *Nature* 2005; 436:214-20.
58. Zhao Y, Ransom JF, Li A, Vedantham V, von Drehle M, Muth AN, et al. Dysregulation of cardiogenesis, cardiac conduction and cell cycle in mice lacking miRNA-1-2. *Cell* 2007; 129:303-17.
59. Kim HK, Lee YS, Sivaprasad U, Malhotra A, Durta A. Muscle-specific microRNA miR-206 promotes muscle differentiation. *J Cell Biol* 2006; 174:677-87.
60. Rosenberg MI, Georges SA, Asawaicharn A, Analau E, Tapscott SJ. MyoD inhibits Fstl1 and Utrn expression by inducing transcription of miR-206. *J Cell Biol* 2006; 175:77-85.
61. Buckingham M, Relaix F. The role of Pax genes in the development of tissues and organs: Pax3 and Pax7 regulate muscle progenitor cell functions. *Annu Rev Cell Dev Biol* 2007; 23:645-73.
62. Crist CG, Montarras D, Pallafacchina G, Rocancourt D, Cumano A, Conway SJ, et al. Muscle stem cell behavior is modified by microRNA-27 regulation of Pax3 expression. *Proc Natl Acad Sci USA* 2009; 106:13383-7.
63. Heremans Y, Van De Castele M, in't Veld P, Gradwohl G, Serup P, Madsen O, et al. Recapitulation of embryonic neuroendocrine differentiation in adult human pancreatic duct cells expressing neurogenin 3. *J Cell Biol* 2002; 159:303-12.
64. Baroukh N, Ravier MA, Loder MK, Hill EV, Bounacer A, Scharfmann R, et al. MicroRNA-124a regulates Foxa2 expression and intracellular signaling in pancreatic beta-cell lines. *J Biol Chem* 2007; 282:19575-88.
65. Fukuda A, Kawaguchi Y, Furuyama K, Kodama S, Horiguchi M, Kuhara T, et al. Ectopic pancreas formation in Hes1-knockout mice reveals plasticity of endodermal progenitors of the gut, bile duct and pancreas. *J Clin Invest* 2006; 116:1484-93.
66. Lynn FC, Skewes-Cox P, Kosaka Y, McManus MT, Harfe BD, German MS. MicroRNA expression is required for pancreatic islet cell genesis in the mouse. *Diabetes* 2007; 56:2938-45.
67. Gradwohl G, Dierich A, LeMeur M, Guillemot F. neurogenin3 is required for the development of the four endocrine cell lineages of the pancreas. *Proc Natl Acad Sci USA* 2000; 97:1607-11.
68. Gu G, Brown JR, Melton DA. Direct lineage tracing reveals the ontogeny of pancreatic cell fates during mouse embryogenesis. *Mech Dev* 2003; 120:35-43.
69. Hara M, Dizon RF, Glick BS, Lee CS, Kaestner KH, Piston DW, et al. Imaging pancreatic beta-cells in the intact pancreas. *Am J Physiol Endocrinol Metab* 2006; 290:1041-7.
70. Jenny M, Uhl C, Roche C, Duluc I, Guillermin V, Guillemot F, et al. Neurogenin3 is differentially required for endocrine cell fate specification in the intestinal and gastric epithelium. *EMBO J* 2002; 21:6338-47.
71. Joglekar MV, Parekh VS, Mehta S, Bhonde RR, Hardikar AA. MicroRNA profiling of developing and regenerating pancreas reveal post-transcriptional regulation of neurogenin3. *Dev Biol* 2007; 311:603-12.
72. Kimura H, Kawasaki H, Taira K. Mouse microRNA-23b regulates expression of Hes1 gene in P19 cells. *Nucleic Acids Symp Ser (Oxf)* 2004; 213-4.
73. Ibayoshi I, Sakamoto M, Yamaguchi M, Mori K, Kageyama R. Essential roles of Notch signaling in maintenance of neural stem cells in developing and adult brains. *J Neurosci* 30:3489-98.
74. Hatakeyama J, Bessho Y, Katoh K, Ookawara S, Fujioka M, Guillemot F, et al. Hes genes regulate size, shape and histogenesis of the nervous system by control of the timing of neural stem cell differentiation. *Development* 2004; 131:5539-50.
75. Bommer GT, Gerin I, Feng Y, Kaczorowski AJ, Kuick R, Love RE, et al. p53-mediated activation of miRNA34 candidate tumor-suppressor genes. *Curr Biol* 2007; 17:1298-307.
76. He L, He X, Lim LP, de Stanchina E, Xuan Z, Liang Y, et al. A microRNA component of the p53 tumour suppressor network. *Nature* 2007; 447:1130-4.
77. Ji Q, Hao X, Zhang M, Tang W, Yang M, Li L, et al. MicroRNA miR-34 inhibits human pancreatic cancer tumor-initiating cells. *PLoS One* 2009; 4:6816.